

## SUMMERTIME MANAGEMENT TIPS

Summertime heat and stress can have a detrimental effect on flock performance. While you cannot control outside weather conditions, you can reduce the effects of summer heat on your flock by following some of these tips to improve bird comfort.

### BROODING

- Set brooder temperatures high enough to cycle. Poults may pile if they cannot find the heat source. If brooders will not cycle due to excessive outside temperatures:
  - Houses with fans - set a fan to run on thermostat
  - Curtain-only houses - crack curtains enough to allow in fresh air
- Turn down any overhead lights to draw poults in under brooders, if required.
- Have fresh water readily available for poults.
- Check poults frequently for piling.

### VENTILATION

- Ensure all fans are in working order, belts are tightened and fan housings are kept free of dust. Clean louvers and vent openings frequently as any dust accumulation will reduce air flow.
- Set and activate alarms. This is especially critical for barns using tunnel and static pressure ventilation.
  - Temperature targets will vary with flock age.
  - Alarms should be tested weekly.
- Properly set thermostats on both fans and curtain machines according to flock age and outside temperature considerations.

- For static-pressure barns, remember the goal is to pull enough static pressure for the air to reach the center of the building while still increasing air volume. *Not all barns are the same width and will have different requirements. Check with your service technician on recommendations for your barn.*
- Set up tunnel-ventilated barns in stages based on outside temperatures and bird requirements.
- For curtain-sided barns, keep screens clean of dust and feathers. Keep perimeter of building clear of tall grass, equipment and any other air-flow obstructions.
- Test any foggers or misters prior to use each summer. Nozzles become clogged and hoses and pipes can become cracked. Systems should be run at service technician recommendations on temperature and timers settings.

### WATER

- Ensure all drinkers are in working order. Adjust drinker height and water depth weekly.
- In barns with nipple drinkers, flush lines daily to keep biofilm buildup to a minimum. Also, check all lines for water to ensure lines are not air locked. Visually check standpipes for proper water pressure.
- Ensure 3-5 PPM free-chlorine reading in last drinker to reduce pathogen load in drinkers. Suggested ORP reading of 650 or more.
- Clean water lines in between flocks to reduce biofilm and other pathogens that can build up during the life of a flock.

## GENERAL TIPS

- Check and run automatic generators weekly, as well as fuel and battery levels.
- Farms equipped with automatic curtain drops for power failure should be hooked up and checked weekly.
- Ensure fly control is in place before warm weather arrives. Flies are known carriers of diseases. Do not wait until there is a problem to begin a program. *Do not assume all products are safe to use with birds present. Check with your service technician on an appropriate program.*
- Consider using electrolytes to reduce stress on birds at key times. Look for electrolyte packs with stabilized vitamin C present. Check with service technician prior to running any product on birds.
  - Excessive heat (>85°F / 29°C) - run electrolytes during daylight hours and fresh water overnight.
  - Moving birds to grow out - run electrolytes the day before move.
  - Loading birds for processing - run electrolytes 24 hours before load.
- Consider using baking soda on younger birds (over the age of 5 weeks) at a rate of 1 lb/1 gal. stock solution during excessive heat (> 90°F / 32°). Check with service technician prior to running any product on birds.
- Collect and dispose of mortality properly; failure to do so can increase the risk of spreading certain diseases. Remember that heat speeds up the decomposition process.
- Keep grass around barns and feed tanks mowed and trimmed. Tall grass and weeds can hinder ventilation, harbor mice, rats, snakes and bees. Rodents can carry disease into your flock. Snakes and bees can be a hazard for growers, service technicians, feed truck drivers and propane delivery drivers.
- Keep end doors and personnel door sealed up to keep skunks and other animals out. These animals can carry disease into your flock.
- Check birds often on days of excessive heat. Make sure all fans and ventilation systems are working. Check water and ensure you have not had any sort of system failure.

The contents of this article are © Aviagen Turkeys and the advice contained herein is given as a guide and may need to be varied to meet a specific customer operation. This advice should not be regarded as a form of guarantee.

